

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

A Newly Identified Member of the Tumor Necrosis Factor Receptor Superfamily with a Wide Tissue Distribution and Involvement in Lymphocyte Activation*

(Received for publication, October 23, 1996, and in revised form, March 25, 1997)

Byoung S. Kwon^{‡§¶}, Kong B. Tan^{§¶}, Jian Ni^{**}, Kwi-Ok-Oh[‡], Zang H Lee^{‡ ¶}, Kack K. Kim[‡], Young-J. Kim[‡], Sa Wang[‡], Reiner Gentz^{**}, Guo-Liang Yu^{**}, Jeremy Harrop[¶], Sally D. Lyn[¶], Carol Silverman[¶], Terence G. Porter[¶], Alem Truneh[¶], and Peter R. Young[¶]

From the [‡]Department of Microbiology and Immunology and Walther Oncology Center, Indiana University School of Medicine, Indianapolis, Indiana 46202, the [¶]Department of Molecular Immunology, King of Prussia, Pennsylvania 19406, and ^{**}Human Genome Sciences, Inc., Rockville, Maryland 20850

The tumor necrosis factor receptor (TNFR) superfamily consists of approximately 10 characterized members of human proteins. We have identified a new member of the TNFR superfamily, TR2, from a search of an expressed sequence tag data base. cDNA cloning and Northern blot hybridization demonstrated multiple mRNA species, of which a 1.7-kilobase form was most abundant. However, TR2 is encoded by a single gene which, maps to chromosome 1p36.22–36.3, in the same region as several other members of the TNFR superfamily. The most abundant TR2 open reading frame encodes a 283-amino acid single transmembrane protein with a 36-residue signal sequence, two perfect and two imperfect TNFR-like cysteine-rich domains, and a short cytoplasmic tail with some similarity to 4-1BB and CD40. TR2 mRNA is expressed in multiple human tissues and cell lines and shows a constitutive and relatively high expression in peripheral blood T cells, B cells, and monocytes. A TR2-Fc fusion protein inhibited a mixed lymphocyte reaction-mediated proliferation suggesting that the receptor and/or its ligand play a role in T cell stimulation.

The members of the tumor necrosis factor receptor (TNFR)^{1/2} nerve growth factor receptor (NGFR) superfamily are characterized by the presence of three to six repeats of a cysteine-rich motif that consists of approximately 30–40 amino acids in the extracellular part of the molecule (1). The crystal structure of TNFR-I complexed with its ligand showed that a cysteine-rich motif (TNFR domain) was composed of three elongated strands of residues held together by a twisted ladder of disulfide bonds

(2). These receptors contain a hinge-like region immediately adjacent to the transmembrane domain, characterized by a lack of cysteine residues and a high proportion of serine, threonine, and proline, which are likely to be glycosylated with O-linked sugars. A cytoplasmic part of these molecules shows limited sequence similarities, a finding that may be the basis for diverse cellular signaling. At present, the members identified from human cells include CD40 (3, 4), 4-1BB (5), OX-40 (6), TNFR-I (7, 8), TNFR-II (9), CD27 (10), Fas (11), NGFR (12), CD30 (13), and LTBR (14). Viral open reading frames encoding soluble TNFRs have also been identified, such as SFV-T2 (9), Va53 (15), G4RG (16), and crmB (17).

Recent studies have shown that these molecules are involved in diverse biological activities such as immunoregulation (18, 19), by regulating cell proliferation (20–22), cell survival (23–25), and cell death (26–28).

Because of their biological significance and the diverse membership of this superfamily, we predicted that there would be further members of the superfamily. By searching an EST data base, we identified a new member of the TNFR superfamily. We report here the initial characterization of the molecule called TR2.

MATERIALS AND METHODS

Identification and Cloning of New Members of the TNFR Superfamily—An EST cDNA data base, obtained from over 500 different cDNA libraries (29, 30), was screened for sequence homology with cysteine-rich motif of the TNFR superfamily, using the blastn and tblastn algorithms (31). One EST was identified in a human T cell line library, which showed significant sequence identity to TNFR-II at the amino acid level. This sequence was used to clone the missing 5' end by RACE (rapid amplification of cDNA ends) using a 5'-RACE ends-ready cDNA from human leukocytes (Clontech, Palo Alto, CA). This sequence matched three further ESTs (HTOBH42, HTOAU65, and HLHA49). Complete sequencing of these and other cDNAs indicated that they contained an identical open reading frame homologous to the TNFR superfamily, and it was named TR2. Analysis of several other ESTs and cDNAs indicated that some cDNAs had additional sequences inserted into the open reading frame identified above and might represent various partially spliced mRNAs.

Cells—The myeloid and B cell lines studied represent cell types at different stages of the differentiation pathway. KG1a and PLB 985 (32, 33) were obtained from Phillip Koeffler (UCLA School of Medicine), BJA-B was from Z Jonak (SmithKline Beecham), and TF 274, a stromal cell line exhibiting osteoblastic features, was generated from the bone marrow of a healthy male donor.² All of the other cell lines were obtained from the American Type Culture Collection (Rockville, MD). Monocytes were prepared by differential centrifugation of peripheral blood mononuclear cells (PBMC) and adhesion to tissue culture dish. CD19⁺, CD4⁺, and CD8⁺ were isolated from PBMC by immunomag-

* This work was supported by National Institutes of Health Grants AI 28175 and AR 40248 (to B. S. K.) and by a postdoctoral fellowship from the American Heart Association, Indiana Affiliate (to S. W.). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

§ The first two authors contributed equally to this work.

¶ To whom correspondence should be addressed. Tel.: 317-274-3950; Fax: 317-274-4090; E-mail: kkwon@sunflower.bio.indiana.edu.

¶ Present address: Dept. of Microbiology, Chosun University School of Dentistry, Kwang Ju 501-759, Korea.

¹ The abbreviations used are: TNFR, tumor necrosis factor receptor; NGFR, nerve growth factor receptor; CHO, Chinese hamster ovary; DAPI, 4,6-diamidino-2-phenylindole; EST, expressed sequence tag; FISH, fluorescein *in situ* hybridization; MLR, mixed lymphocyte reaction; PBMC, peripheral blood mononuclear cell; PHA, phytohemagglutinin; PMA, phorbol myristate acetate; RACE, rapid amplification of cDNA ends; mAb, monoclonal antibody; HVEM, herpesvirus entry mediator.

² K. B. Tan and Z. Jonak, unpublished data.

netic beads (Dynal, Lake Success, NY). Endothelial cells from human coronary artery were purchased from Clonetics (San Diego, CA).

RNA and DNA Blot Hybridization—Total RNA of adult tissues was purchased from CLONTECH or extracted from primary cells and cell lines with TriReagent (Molecular Research Center, Inc., Cincinnati, OH). 5–7.5 μ g of total RNA was fractionated in a 1% agarose gel containing formaldehyde, as described (34), and transferred quantitatively to Zeta-Probe nylon membrane (Bio-Rad) by vacuum blotting. The blots were prehybridized, hybridized with 32 P-labeled *Xho*I/*Eco*RI fragment of TR2 or OX-40 probe, washed under high stringency conditions, and exposed to x-ray films.

High molecular weight human DNA was digested with various restriction enzymes and fractionated in 0.8% agarose gel. The DNA was denatured, neutralized, and transferred to nylon membrane and hybridized to 32 P-labeled TR2 or its variant cDNA.

In Situ Hybridization and FISH Detection—The *in situ* hybridization and FISH detection of TR2 location in human chromosomes were performed as described previously (35, 36). FISH signals and the DAPI banding pattern were recorded separately by taking photographs, and the assignment of the FISH mapping data with chromosomal bands was achieved by superimposing FISH signals with a DAPI-banded chromosome (37).

Production of Recombinant TR2-Fc Fusion Proteins—The 5' portion of the TR2 containing the entire putative open reading frame of extracellular domain was amplified by polymerase chain reaction (38). For correctly oriented cloning, a *Hind*III site on the 5' end of the forward primer and a *Bgl*III site on the 5' end of the reverse primer were created. The Fc portion of human IgG₁ was PCR-amplified from ARH-77 (ATCC) cell RNA and cloned in the *Sma*I site of the pGem7 vector (Promega, Madison, WI). The Fc fragment, including hinge, CH₂, and CH₃ domain sequences, contained a *Bgl*III site at its 5' end and an *Xho*I site at its 3' end. The *Hind*III-*Bgl*III fragment of TR2 cDNA was inserted upstream of human IgG₁-Fc and an in-frame fusion was confirmed by sequencing. The TR2-Fc fragment was released by digesting the plasmid with *Hind*III-*Xho*I and cloned into pcDNA3 expression plasmid.

The TR2-Fc plasmid, linearized with *Pvu*I, was transfected into NIH 3T3 by the calcium phosphate co-precipitation method. After selection in 400 μ g/ml G418, neomycin-resistant colonies were picked and expanded. Enzyme-linked immunosorbent assay with anti-human IgG₁ and Northern analysis with 32 P-labeled TR2 probe were used to select for clones that produce high levels of TR2-Fc in the supernatant. In some experiments, a slightly differently engineered TR2-Fc produced in Chinese hamster ovary (CHO) cells was used. The TR2-Fc was purified by protein G chromatography, and the amino acid sequence of the N terminus was determined by automatic peptide sequencer (ABI).

In Vitro Transcription and Translation—The full-length TR2 cDNA was inserted into *Hind*III-*Xho*I sites of pcDNA 3 vector (Invitrogen, San Diego, CA). TNT-coupled reticulocyte lysate system (Promega) was used to *in vitro* transcribe and translate the TR2 cDNA in pcDNA 3. The 35 S-labeled reaction product was fractionated on a 5–15% gradient SDS-polyacrylamide gel, transferred onto an Immobilon membrane (Millipore, Bedford, MA), and exposed to x-ray film.

Blocking MLR-mediated PBMC Proliferation—PBMC were isolated from three healthy adult volunteers by Ficoll gradient centrifugation at 400 \times *g* for 30 min. PBMCs were recovered, washed in RPMI 1640 (Life Technologies, Inc.) supplemented with 10% fetal bovine serum, 300 μ g/ml L-glutamine, and 50 μ g/ml gentamycin, and adjusted to 1×10^6 cells/ml for two donors and to 2×10^5 cells/ml for the third donor.

Fifty μ l of each cell suspension was added to 96-well (round bottom) plates (Falcon, Franklin Lakes, NJ) together with 50 μ l of TR2-Fc, IL-5R-Fc, anti-CD4 mAb, or control mAb. Plates were incubated at 37 $^{\circ}$ C in 5% CO₂ for 96 h. One μ Ci of [3 H]methylthymidine (ICN Biomedicals, Costa Mesa, CA) was then added for an additional 16 h. Cells were harvested, and radioactivity was counted.

RESULTS AND DISCUSSION

TR2 Is a New Member of the TNFR Superfamily—Fig. 1a shows the amino acid sequence of TR2 deduced from the longest open reading frame of one of the isolated cDNAs (HLHA49). Comparison with other sequenced cDNAs and with ESTs in the data base indicated potential allelic variants that resulted in amino acid changes at positions 17 (either Arg or Lys) and 41 (either Ser or Phe) of the protein sequence.

The open reading frame encodes 283 amino acids with a calculated molecular weight of 30,417. The TR2 protein was expected to be a receptor. Therefore, the potential signal se-




FIG. 1. a, deduced amino acid sequence of TR2. The signal region is underlined. The potential glycosylation sites are underlined with a heavy line. The putative transmembrane region is double underlined. The partial amino acid sequence of recombinant TR2 reads as PALP..., which indicates that the first 36 amino acids constitute a signal sequence. The GenBank™ accession number of this sequence is U81232. b, *in vitro* transcription and translation of TR2 cDNA. TR2 cDNA in pcDNA 3 (lane 1) and reversely oriented TR2 cDNA in pcDNA 3 (lane 2) was used for *in vitro* translation by the TNT-coupled reticulocyte lysate system (Promega). [35 S]Met-labeled translation product was fractionated by a 5–15% gradient SDS-polyacrylamide gel electrophoresis. M indicates molecular size markers, and the arrow indicates TR2 protein band. c, aligned amino acid sequence of extracellular motif of TR2 with other TNFR family members. The amino acid sequence of TR2 was aligned with those of TNFR-I, TNFR-II, CD40, and 4-1BB on the basis of sequence homology and conserved cysteines.

TR2 Maps at 1P36.2-P36.3.—The FISH mapping procedure was applied to localize the TR2 gene to a specific human chromosomal region. The assignment of a hybridization signal to the short arm of chromosome 1 was obtained with the aid of DAPI banding. A total of 10 mitotic figures were photographed, one of which is shown in Fig. 3a. The double fluorescent signals are indicated on the schematic diagram of chromosome 1 as shown in Fig. 3b. This result indicated that the TR2 gene is located on the chromosome 1 region p36.2-p36.3. The TR2




FIG. 3. *a*, combined DAPI banding and *in situ* hybridization. *Panel A*, the position of the TR2 probe is shown by orange dots on a chromosome. *Panel B*, the same chromosome with DAPI banding. *b*, localization of TR2 on human chromosome. A schematic diagram of human chromosome 1 is shown, and the positions at which TR2 hybridization signals were detected are indicated by filled circles.

position is in close proximity with CD30 (42), 4-1BB (43, 44), OX-40 (45), and TNFR-II (46), suggesting that it evolved through a localized gene duplication event. Interestingly, all of these receptors have stimulatory phenotypes in T cells in response to cognate ligand binding, in contrast to Fas and TNFR-I, which stimulate apoptosis. This prompted us to test if TR2 might be involved in lymphocyte stimulation.

TR2-Fc Interferes with MLR-mediated Proliferation of PBMC—To determine the potential involvement of cell surface TR2 with its ligand in lymphocyte proliferation, we examined allogeneic MLR proliferative responses. As shown in Fig. 4, *a* and *b*, when TR2-Fc was added to the culture, a significant reduction of maximal responses was observed ($p < 0.05$). The addition of TR2-Fc at 100 $\mu\text{g/ml}$ inhibited the proliferation up to 53%. No significant inhibition of proliferation was observed with the control IL-5R-Fc. Surprisingly, at high concentrations (10–100 $\mu\text{g/ml}$) IL-5R-Fc was shown to enhance proliferation. The concentrations of TR2-Fc required to inhibit MLR prolif-




FIG. 4. **Inhibition of allogeneic proliferation by TR2-Fc.** *a*, effect of TR2-Fc on a three-way MLR. PBMCs were adjusted to 1×10^6 cells/ml for two donors and to 2×10^6 cells/ml for the third donor. Fifty microliters of each cell suspension was added to 96-well round bottom plates, together with 50 μl of TR2-Fc (○) or with 50 μl of IL-5R-Fc (■). [^3H]Methylthymidine incorporation was measured as an indication of cell proliferation. The error bars represent standard errors. *b*, effect of anti-CD4 mAb on a three-way MLR. PBMCs were adjusted to 1×10^6 cells/ml for two donors and to 2×10^6 cells/ml for the third donor. Fifty microliters of each cell suspension was added to 96-well round bottom plates together with 50 μl of anti-CD4 mAb (○) or with 50 μl of 2B6 mAb (■). [^3H]Methylthymidine incorporation was measured as an indication of cell proliferation. The error bars represent standard errors.

eration (1–100 $\mu\text{g/ml}$) are comparable with those of CD40-Fc required for inhibition in other lymphocyte assays (47–50). An anti-CD4 mAb assayed simultaneously inhibited MLR-mediated proliferation up to 60%, whereas a control anti-IL-5 mAb failed to inhibit the proliferation. It is well known that a major component of the MLR proliferative response is T cell-dependent; hence, it would appear that inhibiting the interaction of TR2 with its ligand prevents optimal T lymphocyte activation and proliferation.

Hence, we have identified an additional member of the TNFR superfamily that either plays a direct role in T cell stimulation or binds to a ligand which can stimulate T cell proliferation through one or more receptors, which may include TR2. We are currently trying to identify this ligand to which TR2 binds to clarify its role.

REFERENCES

1. Mallett, S., and Barclay, A. N. (1991) *Immunol. Today* 12, 220-224
2. Banner, D. W., D'Arcy, A., Jones, W., Gentz, R., Schoenfeld, H.-J., Broger, C., Loetscher, H., and Lesslauer, W. (1993) *Cell* 73, 431-445
3. Banchereau, J., Bazan, F., Blanchard, D., Briere, F., Galizzi, J. P., van Kooten, C., Liu, Y.-J., Rousset, F., and Saeland, S. (1994) *Annu. Rev. Immunol.* 12, 881-922
4. Stamenkovic, I., Clark, E. A., and Seed, B. (1989) *EMBO J.* 8, 1403-1410
5. Kwon, B. S., and Weissman, S. M. (1989) *Proc. Natl. Acad. Sci. U. S. A.* 86, 1963-1967
6. Mallett, S., Fossum, S., and Barclay, A. N. (1990) *EMBO J.* 9, 1063-1068
7. Loetscher, H., Pan, Y.-C., Lohm, H. W., Gentz, R., Brockhaus, M., Tabuchi, H., and Lesslauer, W. (1990) *Cell* 61, 351-359
8. Schall, T. J., Lewis, M., Koller, K. J., Lee, A., Rice, G. C., Wong, G. H., Gatanaga, T., Granger, G. A., Lentz, R., Raab, R., Kohr, W. J., and Goeddel, D. V. (1990) *Cell* 61, 361-370
9. Smith, C. A., Davis, T., Anderson, D., Solam, L., Beckmann, M. P., Jerzy, R., Dower, S. K., Cosman, D., and Goodwin, R. G. (1990) *Science* 248, 1019-1023
10. Van Lier, R. A., Borst, J., Vroom, T. M., Klein, H., Van Mourik, P., Zeijlemaker, W. P., and Melief, C. J. (1987) *J. Immunol.* 139, 1589-1596
11. Itoh, N., Yonehara, S., Ishii, A., Yonehoro, M., Mizushima, S. I., Sameshima, M., Hase, A., Seto, Y., and Nagata, S. (1991) *Cell* 66, 233-243
12. Johnson, D., Lanahan, A., Buck, C. R., Sehgal, A., Morgan, C., Mercer, E., Bothwell, M., and Chao, M. (1986) *Cell* 47, 545-554
13. Durkop, H., Latza, U., Hummel, M., Eitelbach, F., Seed, B., and Stein, H. (1992) *Cell* 68, 421-427
14. Baens, M., Chaffanet, M., Cassiman, J. J., van den Berghe, H., and Marymen, P. (1993) *Genomics* 16, 214-218
15. Smith, C. A., Davis, T., Anderson, D., Solam, L., Beckmann, M. P., Jerzy, R., Dower, S. K., Cosman, D., and Goodwin, R. G. (1990) *Science* 248, 1019-1023
16. Howard, S. T., Chan, Y. S., and Smith, G. L. (1991) *Virology* 180, 633-647
17. Hu, F.-Q., Smith, C. A., and Pickup, D. J. (1994) *Virology* 204, 343-356
18. Smith, G. L. (1993) *J. Gen. Virol.* 74, 1725-1740
19. Armitage, R. J. (1994) *Curr. Opin. Immunol.* 6, 407-413
20. Smith, C. A., Farrah, T., and Goodwin, R. G. (1994) *Cell* 75, 959-962
21. Banchereau, J., de Paoli, P., Valle, A., Garcia, E., and Rousset, P. (1991) *Science* 251, 70-72
22. Pollok, K. E., Kim, Y.-J., Zhou, A., Hurtado, J., Kim, K.-K., Pickard, R. T., and Kwon, B. S. (1993) *J. Immunol.* 150, 771-781
23. Baum, P. R., Gayle, R. B., III, Ramsdell, F., Srinivasan, S., Sorensen, R. A., Watson, M. L., Seldin, M. F., Baker, E., Sutherland, G. R., Clifford, K. N., Alderson, M. R., Goodwin, R. G., and Fanslow, W. C. (1994) *EMBO J.* 13, 3992-4001
24. Grass, H.-J., Boiani, N., Williams, D. E., Armitage, R. J., Smith, C. A., and Goodwin, R. G. (1994) *Blood* 83, 2045-2056
25. Torcia, M., Bracci-Laudiero, L., Lucibello, M., Nencioni, L., Labardi, D., Rubartelli, A., Cozzolino, F., Aloisi, L., and Garaci, E. (1996) *Cell* 85, 345-356
26. Tartaglia, L. A., Ayres, T. M., Wong, G. H., and Goeddel, D. V. (1993) *Cell* 74, 845-853
27. Gillette-Ferguson, I., and Sidman, C. L. (1994) *Eur. J. Immunol.* 24, 1181-1185
28. Krammer, P. H., Behrmann, I., Daniel, P., Dhein, J., and Debatin, K. M. (1994) *Curr. Opin. Immunol.* 6, 279-289
29. Adams, M. D., Kelley, J. M., Gocayne, J. D., Dubnick, M., Polymeropoulos, M. H., Xiao, H., Meril, C. R., Wu, A., Olde, B., and Moreno, R. F. (1991) *Science* 252, 1651-1656
30. Adams, M. D., Dubnick, M., Kerlavage, A. R., Moreno, R. F., Kelley, J. M., Utterback, T. R., Nagle, J. W., Fields, C., and Venter, J. C. (1992) *Nature* 355, 632-634
31. Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990) *J. Mol. Biol.* 215, 403-410
32. Koeffler, H. P., Billing, R., Lusis, A. J., Sparkes, R., and Golde, D. W. (1980) *Blood* 56, 265-273
33. Tucker, K. A., Lilly, M. B., Heck, L., and Rado, T. A. (1987) *Blood* 70, 372-378
34. Sambrook, J., Fritsch, E. F., and Maniatis, J. (1989) *Molecular Cloning: A Laboratory Manual*, Second Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY
35. Heng, H. H. Q., Squire, J., and Tsui, L.-C. (1992) *Proc. Natl. Acad. Sci. U. S. A.* 89, 9509-9513
36. Heng, H. H. Q., Xiao, H., Shi, S.-M., Greenblatt, J., and Tsui, L.-C. (1994) *Hum. Mol. Genet.* 3, 61-64
37. Heng, H. H. Q., and Tsui, L.-C. (1993) *Chromosoma (Berl.)* 102, 325-332
38. Saiki, R. K., Gelfand, D. H., Stoffel, S., Scharf, S. J., Higuchi, R., Horn, G. T., Mullis, K. B., and Erlich, H. A. (1988) *Science* 239, 487-491
39. Montgomery, R. I., Warner, M. S., Lum, B. J., and Spear, P. G. (1996) *Cell* 87, 427-436
40. Banchereau, J., and Rousset, F. (1991) *Nature* 353, 678-679
41. Hurtado, J., Kim, Y.-J., and Kwon, B. S. (1997) *J. Immunol.* 158, 2600-2609
42. Smith, C. A., Gruss, H.-J., Davis-Smith, T., Anderson, D., Farrah, T., Baker, E., Sutherland, G. R., Brannan, C. I., Copeland, N. G., Jenkins, N. A., Grabstein, K. H., Gliniak, B., McAlister, I. B., Fanslow, W., Alderson, M., Falk, B., Gimpel, S., Gillis, S., Din, W. S., Goodwin, R. G., and Armitage, R. J. (1993) *Cell* 73, 1439-1360
43. Kwon, B. S., Kozak, C. A., Kim, K. K., and Pickard, R. T. (1994) *J. Immunol.* 152, 2256-2262
44. Goodwin, R. G., Din, W. S., Davis-Smith, T., Anderson, D. M., Gimpel, S. D., Sato, T. A., Maliszewski, C. R., Brannan, C. L., Copeland, N. G., Jenkins, N. A., Farrah, T., Armitage, R. J., Fanslow, W. C., and Smith, C. A. (1993) *Eur. J. Immunol.* 23, 2631-2641
45. Birkeland, M. L., Copeland, N. G., Gilbert, D. J., Jenkins, N. A., and Barclay, A. N. (1995) *Eur. J. Immunol.* 25, 926-930
46. Baker, E., Chen, L. Z., Smith, C. A., Callen, D. F., Goodwin, R. G., and Sutherland, G. R. (1991) *Cytogenet. Cell Genet.* 57, 117-118
47. Grammer, A. C., Bergman, M. C., Miura, V., Fujita, K., Davis, L. S., and Lipsky, P. E. (1995) *J. Immunol.* 154, 4996-5010
48. Fanslow, W. C., Anderson, D. M., Grabstein, K. H., Clark, E. A., Cosman, D., and Armitage, R. J. (1992) *J. Immunol.* 149, 655-660
49. Grabstein, K. H., Maliszewski, C. R., Shanebeck, K., Sato, T. A., Spriggs, M. K., Fanslow, W. C., and Armitage, R. J. (1993) *J. Immunol.* 150, 3141-3147
50. Noelle, R. J., Roy, M., Shepherd, D. M., Stamenkovic, I., Ledbetter, J. A., and Aruffo, A. (1992) *Proc. Natl. Acad. Sci. U. S. A.* 89, 6550-6554